

Spine Clinics

Cervical Disc Replacement

By Orthopaedics International and Neurosurgery International

Permit No.MICA (P) 149/10/2010

Division of Spine Surgery -
Orthopaedic International



Dr Tan Siah Heng James
MBBS (Singapore) FRCS (Edinburgh)
FRCS (Glasgow),
FRCS (Neurosurgery)
Past Director of Spine Programme NNI ,
Comm Member, East Asia, AO Spine Intl
Member AO Spine International
Member, Clinical Neuroscience Society



Dr Ngian Kite Seng
MBBS (Singapore) FRCS (Edin)
FRCS (Glasgow) FAMS (Ortho)
International Affiliate Member American
Association of Orthopaedic Surgeons
Member, Asian Academy of Minimally Inva-
sive Spinal Surgery (AAMISS)
Member, AOSpine International



Dr Tan Chong Tien
MBBS(Melbourne) FRCS (Edinburgh)
FAMS (Orthopaedics)
Past Director of Spine Service, SGH
Past President Singapore Orthopaedics
Association
Membership: Scoliosis Research Society
Asian Academy of Minimally Invasive Spinal
Surgery

Welcome

Here is our November Issue. We trust you have enjoyed our two previ-
ous issues. We appreciate your positive feedback.

Some of you have asked for an email contact in case of queries. The
author's email is made available should you wish to clarify or require
more information.

Cervical Disc Replacement Surgery

Presentation


Mr C is a 40 year old man who was involved in an accident when he twisted his neck. There was sudden severe neck pain with sharp shooting pain going down his arms. He was treated with medications and physiotherapy for 4 weeks but the pain did not subside and he developed parathesia in his fore-arms.

On examination, he had mild weakness of his biceps and decreased sensation to the C6 dermatome.

Investigations

MRI showed severe prolapsed cervical disc indenting the spinal cord and neuroforamens of C5/6. There was evidence of spinal cord pressure with T2 changes seen on the sagittal views.

Lateral cervical spine flexion and extension x-rays show good movement of the cervical spine. This is important if we are planning to insert and artificial disc.



Axial MRI image showing prolapsed disc at C5/6 compressing the nerve root and deforming the spinal cord.

Treatment

He needed decompression of the spinal cord and nerve roots. Generally, early decompression results in more complete recovery. There are several options to treat this common disorder:

- 1) Anterior cervical disectomy and fusion
- 2) Anterior cervical disectomy and arthroplasty (artificial disc)
- 3) Posterior decompression

We decided to approach this problem via an anterior cervical disectomy and replace the damaged discs with artificial ones.

The advantage to Mr C is that he would not need to lose any cervical mobility, no need to wear a cervical brace post op and no risk of increased adjacent disc disease.

The transverse incision is used in this case. It lends itself to healing very nicely with minimal scarring. Once the damaged and prolapsed disc was removed, the endplates were prepared and the artificial disc inserted. This particular implant does not require too much drilling and is, therefore, safer.

Results

Mr C was admitted for a total of 3 days. He was discharged with no need for any pain medication. The radicular pain and parathesia had fully resolved.



Flexion and extension lateral C-spine x-rays showing the mobility provided by the artificial disc at the operated level



Transverse cervical incisions generally heal very well and can be hidden in the neck creases.

Summary

Cervical disc replacement is a good option for those who require decompression of the cervical spine and would like to maintain their cervical mobility and shorten their recovery process.

Contact email:
dr.james.tan@gmail.com

Orthopaedics International, Neurosurgery International and Sports Medicine International are a group of registered specialist practices comprising 8 orthopaedic surgeons, a neurosurgeon and a sports physician. Operating out of 4 locations within Singapore, we aim to provide patients with comprehensive and professional care for all musculoskeletal, neurosurgical and sports-related conditions. Each specialist brings a range of interests, expertise and sub-specializations to the group, and is also a senior doctor with a minimum of 20 to 30 years of relevant clinical experience behind him. We strongly believe in a team approach, so that every patient of ours will be treated with the highest standards of expertise and care that are available.

Our Locations :

Camden



#03-02 Camden Medical Centre
1 Orchard Boulevard
Singapore 248649
Tel : 6836 9688
Fax : 6836 6869

Mt. Alvernia



#02-33 Mt. Alvernia Medical Centre Blk B
820 Thomson Road
Singapore 574623
Tel : 6352 7678
Fax : 6352 7680

Gleneagles



#02-42 Gleneagles Hospital Annexe Block
6A Napier Road
Singapore 258500
Tel: 6476 7266
Fax: 6476 2066

Mt. Elizabeth



#05-08 Mt Elizabeth Medical Centre
3 Mt Elizabeth
Singapore 228510
Tel. 6737 6386
Fax : 6737 6836

Registered Practices :



Sports Medicine International
a subsidiary of SICRI



Neurosurgery International
a subsidiary of SICRI

www.ortho-intl.com
www.sportsmed-intl.com
www.neuro-intl.com